

# An Investigation of the Kinematic and Microphysical Control of Lightning Rate, Extent and NO<sub>x</sub> Production using DC3 Observations and the NASA Lightning Nitrogen Oxides Model (LNOM)

Lawrence Carey<sup>1</sup>, William Koshak<sup>2</sup>, Harold Peterson<sup>3</sup>,  
Retha Matthee<sup>1</sup> and A. Lamont Bain<sup>1</sup>

<sup>1</sup> Department of Atmospheric Science, University of Alabama in Huntsville (UAH), Huntsville, AL



<sup>2</sup> Earth Science Office, NASA Marshall Space Flight Center (MSFC), Huntsville, AL

<sup>3</sup> Universities Space Research Association (USRA), Huntsville, AL



# Outline

- Deep Convective Clouds and Chemistry (DC<sub>3</sub>) Experiment - Alabama (AL) aircraft case on 21 May 2012
  - Environment
  - Dual-Doppler and dual-polarization radar summary during peak lightning period
  - Lightning properties (rate, extent) vs. radar-inferred kinematic and microphysical characteristics
- NASA Lightning Nitrogen Oxides Model (LNOM)
  - Lightning Segment Altitude Distribution (SAD)
  - Lightning NO<sub>x</sub> (LNO<sub>x</sub>) Production
- LNOM SAD and LNO<sub>x</sub> Production for 21 May 2012 case over AL
  - Comparison to radar



21 May 2012  
 DC3 AL  
 Sounding

## Map of DC3 AL Domain

Low-moderate CAPE, low shear  
 → ordinary multicell storms



## Environmental Conditions

21 May 2012

DC3 AL Aircraft Case  
 2037 UTC UAH RAOB

Convective Parameter	Value
850-700 hPa lapse rate	-3 °C km <sup>-1</sup>
850-500 hPa lapse rate	-6.2 °C km <sup>-1</sup>
SFC-3 km lapse rate	-7.3 °C km <sup>-1</sup>
SBCAPE	785 J kg <sup>-1</sup>
SBCIN	-1 J kg <sup>-1</sup>
DCAPE	702 J kg <sup>-1</sup>
Lifted Index	-2 °C
0-6 km shear	1.2 m s <sup>-1</sup>
0 °C level	3.5 km
-10 °C level	5.5 km
-40 °C level	9.5 km

# LNOM and Radar Analysis Domains







- Initiation approximately around 1940 UTC nearly 80-85 km north of ARMOR
- Multicellular with noticeable cell merger around 2015 UTC
- Peak NA LMA Total Flash Rate  $\sim 5$  flashes  $\text{min}^{-1}$
- Peak NLDN CG Flash Rate  $\sim 1$  flash  $\text{min}^{-1}$
- Peak Vertical Velocity  $\sim 20$   $\text{m s}^{-1}$

# 21 May 2012 (Rapid Intensification, Post Merger 2012-2023 UTC)

## Cell Merger

- Northern updraft congeals with southern updraft
- Continued increase in  $w_{max} \sim 10 \text{ m s}^{-1}$
- Pronounced  $Z_{dr}$  column (5-6 dB)




## Cell Merger

- Strengthening updraft supportive of lofting raindrops upward into MP region for freezing
- Gradient in  $Z_{dr}$  ( $\Delta Z_{dr} \sim 4 \text{ dB} < 2 \text{ km depth}$ ) suggest transition from liquid to frozen particles




# 21 May 2012 (Lightning in Southernmost Updraft 2015 UTC)



## Initial Lightning in "Merged" Cells

- Along gradient of  $Z_h$  and  $Z_{dr}$
- Collocated with  $w_{max} > 8$  m s<sup>-1</sup>
- Storm Scale particle (and thus charge) separation occurred between rimed graupel and ice
- Resultant electrification and flashes



## Z<sub>dr</sub> Column


- High  $Z_h$  (55-60 dBZ) and  $Z_{dr}$  (> 5 dB) into 0 °C to -10 °C layer
- Possible resonance impacts due to high density hail and/or graupel particles
- Persistent  $Z_{dr}$  column is coincident with strong updraft

- LMA flash initiation

..... 0 °C Level  
..... -10 °C Level  
..... -40 °C Level


— W 2, 4, 6, 8, 10 m s<sup>-1</sup>

# 21 May 2012 (Peak Lightning, Post Merger 2023 UTC)





## Peak Lightning

- Peak ~ 5 flashes min<sup>-1</sup>
- Peak likely associated w/ strong 'w' observed at 2015 UTC
- Max 10 dB echo tops above 10 km
- Strong front-to-rear flow (sloped 'w' contours). Efficient at transporting precipitation ice mass aloft for NIC
- Predominately IC flashes aloft with relatively few CG's





Large quantity of graupel given extension of modest  $Z_h$  and low  $Z_{dr}$  into -10 °C to -40 °C layer




Flash Extent = (Convex Hull Flash Area)<sup>1/2</sup>

# Flash Extent (Convex Hull) vs. Flash Count vs. Radar Microphysics/Kinematic




- Flash count correlated with MP precipitation ice and updraft.
- Median flash extent increases as convection pulses
- Largest flashes lag convective pulse but correlated to non-precipitation (anvil) ice
- Flash count and extent opposed; most large flashes when flash rate low (e.g., [Bruning and MacGorman 2013](#))






# NASA Lightning Nitrogen Oxides Model (LNOM)




Koshak et al. (2013)

- LNOM run in Lagrangian (i.e., storm following cylinder) mode for multi-cell cluster sampled by DC3 aircraft on 21 May 2012 over AL
- Subjective radar- and LMA-based definition of multicell cluster, shown earlier
- Variable LNOM cylinder radius size and location that change each ARMOR radar volume time


# NASA LNOm Lightning Segment Altitude Distribution (SAD)



# Max dBZ



## Ground Flashes




# NASA LNOm Lightning NO<sub>x</sub> (LNO<sub>x</sub>) Production


- Cloud flashes dominate SAD
- Ground SAD  $\leq$  Cloud SAD, especially aloft
- Yet, LNO<sub>x</sub> production from Ground Flash significant fraction of overall LNO<sub>x</sub> production, especially but not exclusively at low levels

# Storm Integrated ( $\approx$ 1-hour) LNO<sub>x</sub> Profiles 21 May 2012 DC3 AL Aircraft Cluster

## SAD All Flashes




## LNO<sub>x</sub> Production All Flashes



# LNOM Flash Extent [ $\Sigma(\text{SAD})$ ] vs. Convex Hull Length vs. Radar


SAD: “connect-the-dots”

Convex Hull: “polygon wrapper”



LNOM Flash

$$\rho(\text{Rate, Extent}) = 0.90$$





Radar Parameter	Flash Rate (min <sup>-1</sup> )	Flash Extent Rate (km min <sup>-1</sup> )
Graupel Echo Volume	$\rho = 0.79$	$\rho = 0.61$
Precipitation Ice Mass	$\rho = 0.78$	$\rho = 0.55$
Updraft Echo Volume	$\rho = 0.76$	$\rho = 0.61$
Maximum Updraft	$\rho = 0.60$	$\rho = 0.41$



# LNOM LNO<sub>x</sub> Production



- Cloud LNO<sub>x</sub> production highly correlated to total cloud flash extent;
- Ground LNO<sub>x</sub> production also correlated to total ground flash extent
- Ground LNO<sub>x</sub> governed in part by other CG flash parameters (e.g., peak current) in LNOM
- Radar microphysical parameters (e.g., graupel volume) somewhat correlated ( $\rho=0.47$ ) to Total LNO<sub>x</sub> Production
  - LNO<sub>x</sub> lags graupel ( $\rho_{lag}=0.64$ )



# Summary

- Dual-Doppler, dual-polarization radar, LMA and LNOM study of the 21 May 2012 DC3 aircraft case over Alabama
- Coalescence-freezing, modest convective updrafts and subsequent graupel growth drives lightning production
- Total lightning flash rate well correlated to kinematic (e.g., updraft volume) and microphysical properties (e.g., graupel volume) inferred from radar, as in past studies
- To a somewhat lesser extent, LNOM flash extent and LNO<sub>x</sub> production also correlated to radar properties
- LNOM SAD (“connect the dots”) flash extent well correlated to convex hull length scale of flash
- Flash count and extent opposed; largest extent flashes lag the convective generator.

# 21 May 2012 Development Phase (1945-2001 UTC)



## $Z_{dr}$ Column


- $Z_{dr} \sim 2-3$  dB
- $Z_h \sim 50-55$  dBZ
- raindrops

## Warm Rain Coalescence

- $Z_{dr} \sim 3-4$  dB
- $Z_h \sim 40-50$  dBZ
- raindrops

## MP Hydrometeor Region

- $Z_{dr}$  0-1 dB
- $Z_h \sim 40-45$  dBZ
- $w_{max} \sim 2-4$  m  $s^{-1}$
- Depressed values of  $\rho_{hv}$  (not shown) suggest mix of freezing/frozen raindrops, graupel/small hail



## 6 m $s^{-1}$ Updraft



- Peaks into 0 °C to -10 °C layer
- Sufficient to loft liquid drops into MP zone
- Favors freezing of rain drops with time

# 21 May 2012 (Electrification of Northernmost Updraft 2004 UTC)

Sufficiently strong vertical motion to loft rain into mixed phase (MP) where freezing by 2001 UTC, likely resulted in the northernmost updraft producing 3 LMA flashes by 2004 UTC.

## Northern Updraft

- $Z_{dr} < 1$  dB
- $Z_h \sim 45-50$  dBZ
- Likely Graupel/Small Hail (PID Confirmed)



- LMA flash initiation

## First Lightning

- First flashes are associated with northern updraft

## Northern Updraft

- Short lived & decayed
- $Z_h$  decrease to 50-55 dBZ
- $W_{max} \sim 2-4$  m s<sup>-1</sup>

## Southern Updraft


- Strengthens
- $Z_h$  decrease to 55 dBZ
- $W_{max} \sim 5-10$  m s<sup>-1</sup>



# 21 May 2012 (Decay and Dissipation Stages 2106 UTC)

## Gradual Decay

- Anvil (rich with ice crystals) extends nearly 40-50 km north of previous convection, but little to no graupel
- $W_{\max} < 4 \text{ m s}^{-1}$
- TLF<sub>R</sub> decreases to  $\sim 1 \text{ flash min}^{-1}$
- More extensive flashes in anvil region



## New Convection

- Classic multicell evolution with additional cells developing along southwestward moving gust front

